

Intertwining Operator Realization of the AdS/CFT Correspondence

V.K. Dobrev*

Arnold Sommerfeld Institute for Mathematical Physics
Technical University Clausthal
Leibnizstr. 10, 38678 Clausthal-Zellerfeld, Germany
asvd@rz.tu-clausthal.de

Abstract

We give a group-theoretic interpretation of the AdS/CFT correspondence as relation of representation equivalence between representations of the conformal group describing the bulk AdS fields ϕ , their boundary fields ϕ_0 and the coupled to the latter boundary conformal operators $\mathcal{O}$. We use two kinds of equivalences. The first kind is equivalence between the representations describing the bulk fields and the boundary fields and it is established here. The second kind is the equivalence between conjugated conformal representations related by Weyl reflection, e.g., the coupled fields ϕ_0 and $\mathcal{O}$. Operators realizing the first kind of equivalence for special cases were actually given by Witten and others - here they are constructed in a more general setting from the requirement that they are intertwining operators. The intertwining operators realizing the second kind of equivalence are provided by the standard conformal two-point functions. Using both equivalences we find that the bulk field has in fact two boundary fields, namely, the coupled fields ϕ_0 and $\mathcal{O}$, the limits being governed by the corresponding conjugated conformal weights $d-\Delta$ and Δ . Thus, from the viewpoint of the bulk-boundary correspondence the coupled fields ϕ_0 and $\mathcal{O}$ are generically on an equal footing.

Our setting is more general since our bulk fields are described by representations of the Euclidean conformal group, i.e., the de Sitter group $G = SO(d+1, 1)$, induced from representations τ of the maximal compact subgroup $SO(d+1)$ of G . From these large reducible representations we can single out representations which are equivalent to conformal boundary representations labelled by the conformal weight and by arbitrary representations μ of the Euclidean Lorentz group $M = SO(d)$, such that μ is contained in the restriction of τ to M . Thus, our boundary $\leftrightarrow$ bulk operators can be compared with those in the literature only when for a fixed μ we consider a 'minimal' representation $\tau = \tau(\mu)$ containing μ . We also relate the boundary $\rightarrow$ bulk normalization constant to the Plancherel measure for G .

* Permanent address: Bulgarian Academy of Sciences, Institute of Nuclear Research and Nuclear Energy, 72 Tsarigradsko Chaussee, 1784 Sofia, Bulgaria.

1. Introduction

Recently there was renewed interest in (super)conformal field theories in arbitrary dimensions. This happened after the remarkable proposal in [1], according to which the large N limit of a conformally invariant theory in d dimensions is governed by supergravity (and string theory) on $d + 1$ -dimensional AdS space (often called AdS_{d+1}) times a compact manifold. Actually the possible relation of field theory on AdS_{d+1} to field theory on $\mathcal{M}_d$ has been a subject of long interest, cf., e.g., [2–6], and also [7–9] for discussions motivated by recent developments, and additional references. The proposal of [1] was elaborated in [10] and [11] where was proposed a precise correspondence between conformal field theory observables and those of supergravity: correlation functions in conformal field theory are given by the dependence of the supergravity action on the asymptotic behavior at infinity. More explicitly, a conformal field $\mathcal{O}$ corresponds to an AdS field ϕ when there exists a conformal invariant coupling $\int \phi_0 \mathcal{O}$ where ϕ_0 is the value of ϕ at the boundary of AdS_{d+1} [11]. Furthermore, the dimension Δ of the operator $\mathcal{O}$ is given by the mass of the particle described by ϕ in supergravity [11]. Also the spectrum of Kaluza-Klein excitations of $AdS_5 \times S^5$, as computed in [12,13], can be matched precisely with certain operators of the $N = 4$ super Yang-Mills theory in four dimensions [11]. After these initial papers there was an explosion of related research of which of interest to us are two aspects: 1) calculation of conformal correlators from AdS (super)gravity, cf., e.g., [14–33]; 2) matching of gravity and string spectra with conformal theories, cf., e.g., [34–47].

One of the main features furnishing this correspondence is that the boundary $\mathcal{M}_d$ of AdS_{d+1} is in fact a copy of d -dimensional Minkowski space (with a cone added at infinity); the symmetry group $SO(d, 2)$ of AdS_{d+1} acts on $\mathcal{M}_d$ as the conformal group. The fact that $SO(d, 2)$ acts on AdS_{d+1} as a group of ordinary symmetries and on $\mathcal{M}_d$ as a group of conformal symmetries means that there are two ways to get a physical theory with $SO(d, 2)$ symmetry: in a relativistic field theory (with or without gravity) on AdS_{d+1} , or in a conformal field theory on $\mathcal{M}_d$.

The main aim of this paper is to give a group-theoretic interpretation of the above correspondence. In fact such an interpretation is partially present in [48] for the $d = 3$ Euclidean version of the AdS/CFT correspondence in the context of the construction of discrete series representations of the group $SO(4, 1)$ involving symmetric traceless tensors of arbitrary nonzero spin.

In short the essence of our interpretation is that the above correspondence is a relation of *representation equivalence* between the representations describing the fields ϕ , ϕ_0 and $\mathcal{O}$. There are actually two kinds of equivalences. The first kind is new (besides the example from [48] mentioned above) and is proved here - it is between the representations describing the bulk fields and the boundary fields. The second kind is well known - it is the equivalence between boundary conformal representations which are related by Weyl reflections, the representations here being the coupled fields ϕ_0 and $\mathcal{O}$.

Our interpretation means that the operators relating these fields are *intertwining operators* between (partially) equivalent representations. Operators giving the first kind of equivalence for special cases were actually given in [11,14,17–19,25–27] - here they are

constructed in a more general setting from the requirement that they are intertwining operators. The operators giving the second kind of equivalence are provided by the standard conformal two-point functions (and the latter intertwining property was known long time ago, cf. [49], also [48]). Using both equivalences we find that the bulk field has naturally *two* boundary fields, namely, the coupled fields ϕ_0 and $\mathcal{O}$, the limits being governed by the corresponding conjugated conformal weights $d - \Delta$ and Δ . Thus, we notice that from the point of view of the bulk-to-boundary correspondence the coupled fields ϕ_0 and $\mathcal{O}$ are generically *on an equal footing*.

As mentioned, *our setting is more general*. In order to be more specific we consider here the Euclidean version of the AdS/CFT correspondence. In this case the conformal group is the *de Sitter group* $G = SO(d + 1, 1)$. Our bulk fields are obtained from representations of G induced from representations τ of the maximal compact subgroup $K = SO(d + 1)$ of G . From these large reducible representations we can single out representations which are equivalent to conformal boundary representations labelled by arbitrary representations μ of the Euclidean Lorentz group $M = SO(d)$, such that μ is contained in the restriction of τ to M . Thus, our boundary $\leftrightarrow$ bulk operators can be compared with those in the literature only when for a fixed μ we consider a 'minimal' representation $\tau = \tau(\mu)$ containing μ . We also relate the boundary $\rightarrow$ bulk normalization constant to the Plancherel measure for G .

On the AdS side the representations are realized on *de Sitter space* $\mathcal{S}$ (the Euclidean counterpart of AdS space), which is isomorphic to the coset G/K . What is also very essential for our approach is that $\mathcal{S}$ is isomorphic (via the Iwasawa decomposition) also to the solvable product group $\tilde{N}A$, where $\tilde{N}$ is the abelian group of Euclidean translations (isomorphic to $\mathbb{R}^d$), A is the one-dimensional dilatation group. The isomorphism $\mathcal{S} \cong \tilde{N}A$ and related ones are explicated in Section 2. On the conformal side the representations are realized on $\tilde{N}$, and we use also the fact that the latter is locally isomorphic (via the Bruhat decomposition) to the coset G/MAN , (where N is the group of special conformal transformations). These representations, called elementary representations (ERs), are introduced in Section 3. The representations on de Sitter space are introduced in Section 4. Then in Section 5 we give the bulk-to-boundary and boundary-to-bulk intertwining operators and discuss the difference between equivalence and partial equivalence. There we display the second limit of the bulk fields and we derive some further intertwining relations. From the latter we derive the relation to the Plancherel measure for G . We illustrate the intertwining relations by a commutative diagram. Section 6 contains some more comments and outlook.

2. de Sitter space from Iwasawa decomposition

As we mentioned the relation between the two pictures uses the fact that $\mathbb{R}^d$ is easily identified within the $d + 1$ -dimensional de Sitter space. Indeed, de Sitter space may be parametrized as:

$$\xi_{d+2}^2 - \sum_{\alpha=1}^{d+1} \xi_{\alpha}^2 = 1, \quad \xi_{d+2} \geq 1 \quad (2.1)$$

and the first d of the ξ_{α} may be taken as coordinates on $\mathbb{R}^d$.

The group-theoretic interpretation of this relation is present in [48] using the so-called *Iwasawa* decomposition $G = \tilde{N}AK$.¹ This is a global decomposition, i.e., each element g of G can be decomposed as the product of the corresponding matrices: $g = \tilde{n}ak$, with $\tilde{n} \in \tilde{N}$, $a \in A$, $k \in K$. To be explicit we use the following defining relation of G :

$$G = \{g \in GL(d+2, \mathbb{R}) \mid {}^t g \eta g = \eta \doteq \text{diag}(-1, \dots, -1, 1), \det g = 1, g_{d+2, d+2} \geq 1\} \quad (2.2)$$

which is the identity component of $O(d+1, 1)$, (${}^t g$ is the transposed of g). Then we have the following matrix representations of the necessary subgroup elements (cf. formulae (1.20a), (1.21), (1.23), of [48], with $2h$ replaced here by d):

$$\tilde{n} = \tilde{n}_x = \begin{pmatrix} \delta_{ij} & -x_i & x_i \\ x_j & 1 - \frac{1}{2}x^2 & \frac{1}{2}x^2 \\ x_j & -\frac{1}{2}x^2 & 1 + \frac{1}{2}x^2 \end{pmatrix} \in \tilde{N}, \quad x \in \mathbb{R}^d, \quad x^2 \equiv \sum_{i=1}^d x_i^2 \quad (2.3a)$$

$$a = \begin{pmatrix} \delta_{ij} & 0 & 0 \\ 0 & \frac{|a|^2+1}{2|a|} & \frac{|a|^2-1}{2|a|} \\ 0 & \frac{|a|^2-1}{2|a|} & \frac{|a|^2+1}{2|a|} \end{pmatrix} \in A, \quad |a| > 0 \quad (2.3b)$$

$$k = \begin{pmatrix} k_{ij} & k_{i, d+1} & 0 \\ k_{d+1, j} & k_{d+1, d+1} & 0 \\ 0 & 0 & 1 \end{pmatrix} \in K, \quad (k_{\alpha\beta}) \in SO(d+1) \quad (2.3c)$$

Writing $g = \tilde{n}ak$ one may determine the factors $\tilde{n}, a, k$ in terms of the matrix elements of g . We use this for the elements of the last column of g , which actually parametrize the de Sitter space, i.e.,

$$\xi_A = g_{A, d+2}, \quad A = 1, \dots, d+2. \quad (2.4)$$

Indeed, substituting (2.4) in (2.1) we recover the $(d+2, d+2)$ -element of the defining relation (2.2), i.e.,

$$\begin{aligned} ({}^t g \eta g)_{d+2, d+2} &= \eta_{d+2, d+2} \implies \\ g_{d+2, d+2}^2 - \sum_{a=1}^{d+1} g_{a, d+2}^2 &= 1 \end{aligned} \quad (2.5)$$

¹ There are several versions of the Iwasawa decomposition, e.g., one may use the group N instead of $\tilde{N}$, and one may take different order of the three factors involved. The choice of version is a matter of convenience.

Now in terms of the parameters in (2.3) we get for the elements of the last column of g , resp., for the parameters of de Sitter space :

$$\begin{aligned} g_{i,d+2} &= \xi_i = \frac{1}{|a|} x_i \\ g_{d+1,d+2} &= \xi_{d+1} = \frac{|a|^2 - 1 + x^2}{2|a|} \\ g_{d+2,d+2} &= \xi_{d+2} = \frac{|a|^2 + 1 + x^2}{2|a|} \geq 1 \end{aligned} \quad (2.6)$$

Notice that the only the $d+1$ parameters of the matrices $\tilde{n}_x, a$ of the Iwasawa decomposition (cf. (2.3a, b)) are involved. Solving (2.6) we obtain for the latter parameters:

$$\begin{aligned} x_i &= \frac{g_{i,d+2}}{g_{d+2,d+2} - g_{d+1,d+2}} = \frac{\xi_i}{\xi_{d+2} - \xi_{d+1}} \\ |a| &= \frac{1}{g_{d+2,d+2} - g_{d+1,d+2}} = \frac{1}{\xi_{d+2} - \xi_{d+1}} > 0 \end{aligned} \quad (2.7)$$

(The last condition follows from $g_{d+2,d+2} \geq 1$.) From (2.6) we get also the consistency condition:

$$\frac{|a|^2 + x^2}{|a|} = g_{d+2,d+2} + g_{d+1,d+2} = \xi_{d+2} + \xi_{d+1} \quad (2.8)$$

Indeed, inserting (2.7) in (2.8) we recover (2.1) and (2.5).

- Thus, in (2.6) and (2.7) we have the mentioned correspondence between de Sitter space $\mathcal{S}$ and the (coset) solvable subgroup $S = \tilde{N}A \cong G/K \cong \mathbb{R}^d \times \mathbb{R}_{>0}$ of the de Sitter group G . In addition, this explicates the group-theoretical interpretation of Euclidean space-time $\mathbb{R}^d$ as the abelian subgroup $\tilde{N}$ of the solvable subgroup S , and the topological interpretation of $\mathbb{R}^d$ as the boundary of $\mathbb{R}^d \times \mathbb{R}_{>0}$ for $|a| \rightarrow 0$.

Remark: Note that for d even some expressions are simpler if we work with the extended de Sitter group:

$$G' \doteq \{g \in GL(d+2, \mathbb{R}) \mid {}^t g \eta g = \eta, \quad g_{d+2,d+2} \geq 1\} \quad (2.9)$$

which includes reflections of the first $d+1$ axes. Then $K \rightarrow K' = O(d+1)$, $M \rightarrow M' = O(d)$, but the de Sitter space $\mathcal{S}$ and the results of this Section are not changed. $\diamond$

3. Conformal field theory representations

The representations used in conformal field theory are called (in the representation theory of semisimple Lie groups) generalized principal series representations (cf. [50]). In [48], [51], [52] they were called *elementary representations* (ERs). They are obtained by induction from the subgroup $P = MAN$, (P is called a parabolic subgroup of G). The induction is from unitary irreps of $M = SO(d)$, from arbitrary (non-unitary) characters of A , and trivially from N . There are several realizations of these representations. We give now the so-called *noncompact picture* of the ERs - it is the one actually used in physics.

The representation space of these induced representations consists of smooth functions on $\mathbb{R}^d$ with values in the corresponding finite-dimensional representation space of M , i.e.:

$$C_\chi = \{f \in C^\infty(\mathbb{R}^d, V_\mu)\} \quad (3.1)$$

where $\chi = [\mu, \Delta]$, Δ is the conformal weight, μ is a unitary irrep of M , V_μ is the finite-dimensional representation space of μ . In addition, these functions have special asymptotic expansion as $x \rightarrow \infty$. The leading term of this expansion is $f(x) \sim \frac{1}{(x^2)^\Delta} f_0$, (for more details of this expansion we refer to [48], [51], [52]). The representation T^χ acts in C_χ by:

$$(T^\chi(g)f)(x) = |a|^{-\Delta} \cdot D^\mu(m) f(x') \quad (3.2)$$

where the nonglobal Bruhat decomposition $g = \tilde{n}man$ is used:²

$$g^{-1}\tilde{n}_x = \tilde{n}_{x'}m^{-1}a^{-1}n^{-1}, \quad g \in G, \tilde{n}_x, \tilde{n}_{x'} \in \tilde{N}, m \in M, a \in A, n \in N \quad (3.3)$$

$D^\mu(m)$ is the representation matrix of μ in V_μ .³

The ERs are generically irreducible both operatorially (in the sense of Schur's Lemma) and topologically (meaning nonexistence of nontrivial (closed) invariant subspaces). However, most of the physically relevant examples are when the ERs are topologically reducible and indecomposable. In particular, such are the representations describing gauge fields, cf., e.g., [53–56].

The importance of the elementary representations comes also from the remarkable result of Langlands-Knapp-Zuckerman [57], [58] stating that every irreducible admissible representation of a real connected semisimple Lie group G with finite centre is equivalent to

² For the cases with measure zero for which $g^{-1}\tilde{n}_x$ does not have a Bruhat decomposition of the form $\tilde{n}man$ the action is defined separately, and the passage from (3.2) to these special cases is ensured to be smooth by the asymptotic properties mentioned above. Further, we may omit such measure-zero exceptions from the formulae - in a rigorous exposition all of them are taken care of, cf. [48], [51], [52].

³ One may interpret C_χ also as a space of smooth sections of the homogeneous vector bundle with base space G/MAN and fibre V_μ .

a subrepresentation of an elementary representation of G .⁴ To obtain a subrepresentation of a topologically reducible ER one has to solve certain invariant differential equations, cf. [48], [51], [52].

Note that the representation data given by $\chi = [\mu, \Delta]$ fixes also the value of the Casimir operators $\mathcal{C}_i$ in the ER C_χ , independently of the latter reducibility. For later use we write:

$$\mathcal{C}_i(\{X\})f(x) = \lambda_i(\mu, \Delta)f(x), \quad i = 1, \dots, \text{rank } G = [\frac{d}{2}] + 1, \quad (3.4)$$

where $\{X\}$ denotes symbolically the generators of the Lie algebra $\mathcal{G}$ of G , and the action of $X \in \mathcal{G}$ is given by the infinitesimal version of (3.2):

$$(Xf)(x) \doteq \frac{\partial}{\partial t} (T^X(\exp tX)f)(x)|_{t=0} \quad (3.5)$$

applying the Bruhat decomposition to $\exp(-tX)\tilde{n}_x$.

Next, we would like to recall the general expression of the conformal two-point function $G_\chi(x_1 - x_2)$ (for special cases cf. [59–62], for the general formula with special stress on the role of the conformal inversion, cf. [49], also [48]):

$$G_\chi(x) = \frac{\gamma_\chi}{(x^2)^\Delta} D^\mu(r(x)) \quad (3.6)$$

$$r(x) = \begin{pmatrix} \tilde{r}(x) & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \in M, \quad \tilde{r}(x) = \left(\frac{2}{x^2} x_i x_j - \delta_{ij} \right)$$

where γ_χ is an arbitrary constant for the moment. (Note that for d even $r(x) \in O(d)$, so we work with G' , cf. (2.9).)

Finally, we note the intertwining property of $G_\chi(x)$. Namely, let $\tilde{\chi}$ be the representation conjugated to χ by Weyl reflection, i.e., by the nontrivial element of the two-element restricted Weyl group $W(G, A)$ [48]. Then we have:

$$\tilde{\chi} \doteq [\tilde{\mu}, d - \Delta], \quad \text{for } \chi = [\mu, \Delta], \quad (3.7)$$

where $\tilde{\mu}$ is the *mirror image* representation of μ . (For d odd $\tilde{\mu} \cong \mu$, while for d even $\tilde{\mu}$ may be obtained from μ by exchanging the representation labels of the two distinguished Dynkin nodes of $SO(d)$.) Then there is the following intertwining operator [49], [48]:

$$G_\chi : C_{\tilde{\chi}} \longrightarrow C_\chi, \quad T^X(g) \circ G_\chi = G_\chi \circ T^{\tilde{X}}(g), \quad \forall g, \quad (3.8a)$$

$$(G_\chi f)(x_1) \doteq \int G_\chi(x_1 - x_2) f(x_2) dx_2. \quad (3.8b)$$

⁴ *Subrepresentations* are irreducible representations realized on invariant subspaces of the ER spaces (in particular, the irreducible ERs themselves). The admissibility condition is fulfilled in the physically interesting examples.

($dx \equiv d^d x$) which means that the representations are partially equivalent. Note that because of this equivalence the values of all Casimirs coincide:

$$\lambda_i(\tilde{\mu}, d - \Delta) = \lambda_i(\mu, \Delta), \quad \forall i. \quad (3.9)$$

Note, that at generic points the representations are equivalent, namely, one has [48], [51]:

$$G_\chi G_{\tilde{\chi}} = \mathbf{1}_\chi, \quad G_{\tilde{\chi}} G_\chi = \mathbf{1}_{\tilde{\chi}} \quad (3.10)$$

This may be used to fix the constant γ_χ .

From the point of view of the AdS/CFT correspondence the importance of the pair $\chi, \tilde{\chi}$ is in the fact that the corresponding fields have conformally invariant coupling through the standard bilinear form:

$$\langle \phi_0, \mathcal{O} \rangle \doteq \int dx \langle \phi_0(x), \mathcal{O}(x) \rangle_\mu, \quad \phi_0 \in C_{\tilde{\chi}}, \quad \mathcal{O} \in C_\chi, \quad (3.11)$$

where $\langle \cdot, \cdot \rangle_\mu$ is the standard pairing between μ and $\tilde{\mu}$. (Note that if ϕ_0 determines a p -form, then $\mathcal{O}(x)$ determines a $(d - p)$ -form.)

4. Representations on de Sitter space

In the previous section we discussed representations on $\mathbb{R}^d \cong \tilde{N}$ induced from the parabolic subgroup MAN which is natural since the abelian subgroup $\tilde{N}$ is locally isomorphic to the factor space G/MAN (via the Bruhat decomposition). Similarly, it is natural to discuss representations on de Sitter space $\mathcal{S} \cong \tilde{N}A$ which are induced from the maximal compact subgroup $K = SO(d + 1)$ since the solvable group $\tilde{N}A$ is isomorphic to the factor space G/K (via the Iwasawa decomposition). Namely, we consider the representation space:

$$\hat{C}_\tau = \{\phi \in C^\infty(\mathbb{R}^d \times \mathbb{R}_{>0}, U_\tau)\} \quad (4.1)$$

where τ is an arbitrary unitary irrep of K , U_τ is the finite-dimensional representation space of τ , with representation action:

$$(\hat{T}^\tau(g)\phi)(x, |a|) = \hat{D}^\tau(k) \phi(x', |a'|) \quad (4.2)$$

where the Iwasawa decomposition is used:

$$g^{-1} \tilde{n}_x a = \tilde{n}_{x'} a' k^{-1}, \quad g \in G, k \in K, \tilde{n}_x, \tilde{n}_{x'} \in \tilde{N}, a, a' \in A \quad (4.3)$$

and $\hat{D}^\tau(k)$ is the representation matrix of τ in U_τ . However, unlike the ERs, these representations are reducible, and to single out an irrep equivalent, say, a subrepresentation of an ER, one has to look for solutions of the eigenvalue problem related to the Casimir operators. This procedure is actually well understood and used in the construction of the discrete series of unitary representations, cf. [63], [64], (also [48] for $d = 3$).

In the actual implementation of (4.2) it is convenient to use the unique decomposition:

$$k = m(k)k_f, \quad m(k) = \begin{pmatrix} \tilde{m}(k) & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \in M, \quad k_f = \begin{pmatrix} \tilde{k}_f & 0 \\ 0 & 1 \end{pmatrix} \in K \quad (4.4)$$

representing the decomposition of K into its subgroup M and the coset $K/M : K \cong M K/M$. Explicitly, using the parametrization of k in (2.3c), we have (for $k_{d+1,d+1} \neq -1$):

$$\begin{aligned} \tilde{m}(k) &= \begin{pmatrix} k_{ij} - \frac{1}{1+k_{d+1,d+1}} k_{i,d+1} k_{d+1,j} \\ \delta_{ij} - \frac{1}{1+k_{d+1,d+1}} k_{d+1,i} k_{d+1,j} & -k_{d+1,i} \\ k_{d+1,j} & k_{d+1,d+1} \end{pmatrix} = \\ \tilde{k}_f &= \begin{pmatrix} \delta_{ij} - \frac{2}{1+x^2} x_i x_j & -\frac{2}{1+x^2} x_i \\ \frac{2}{1+x^2} x_j & \frac{1-x^2}{1+x^2} \end{pmatrix} \doteq \tilde{k}_x \\ x \in \mathbb{R}^d, \quad x_i &= \frac{1}{1+k_{d+1,d+1}} k_{d+1,i}, \quad x^2 = \frac{1-k_{d+1,d+1}}{1+k_{d+1,d+1}} \end{aligned} \quad (4.5)$$

Note that $k_x \doteq \begin{pmatrix} \tilde{k}_x & 0 \\ 0 & 1 \end{pmatrix}$ appeared in (1.30a) of [48].⁵

Further, we would like to extract from $\hat{C}_\tau$ a representation that may be equivalent to C_χ , $\chi = [\mu, \Delta]$. The first condition for this is that the M -representation μ is contained in the restriction of the K -representation τ to M . Another condition is that the two representations would have the same Casimir values $\lambda_i(\mu, \Delta)$. Having in mind the degeneracy of Casimir values for partially equivalent representations (e.g., (3.9)) we add also the appropriate asymptotic condition. Furthermore, from now on we shall suppose that Δ is real. Thus, we shall use the representations:

$$\begin{aligned} \hat{C}_\chi^\tau &= \{ \phi \in \hat{C}_\tau : \mathcal{C}_i(\{\hat{X}\}) \phi(x, |a|) = \lambda_i(\mu, \Delta) \phi(x, |a|), \quad \forall i, \quad \mu \in \tau|_M, \\ &\quad \phi(x, |a|) \sim |a|^\Delta \varphi(x) \text{ for } |a| \rightarrow 0 \} \end{aligned} \quad (4.6)$$

where $\{\hat{X}\}$ denotes symbolically the generators of the Lie algebra $\mathcal{G}$ with the action $\hat{X}$ of $X \in \mathcal{G}$ given by the infinitesimal version of (4.2):

$$(\hat{X} \phi)(x, |a|) \doteq \frac{\partial}{\partial t} (\hat{T}^\tau(\exp tX) \phi)(x, |a|)|_{t=0} \quad (4.7)$$

applying the Iwasawa decomposition to $\exp(-tX) \tilde{n}_x a$. Certainly, the Casimirs $\mathcal{C}_i(\{\hat{X}\})$ with (4.7) substituted are differential operators and the elements of $\hat{C}_\chi$ are solutions of the equations above.

⁵ The matrices k_x realize the (local) isomorphisms: $\mathbb{R}^d \cong \tilde{N} \stackrel{\text{loc}}{\cong} K/M \cong G/MAN$ (using for the last isomorphism the Iwasawa decomposition in the version $G = KAN$).

Remark: Note that generically the functions in (4.6) have also a second limit with $\Delta \rightarrow d - \Delta$:

$$\tilde{\varphi}(x) = \lim_{|a| \rightarrow 0} |a|^{\Delta-d} \phi(x, |a|) \quad (4.8)$$

which will appear as a consequence of the formalism. With this we shall establish - for generic representations - the following important relation:

$$\hat{C}_\chi^\tau = \hat{C}_{\tilde{\chi}}^\tau, \quad \chi = [\mu, \Delta], \quad \tilde{\chi} = [\tilde{\mu}, d - \Delta], \quad (4.9)$$

for which besides (4.8), we use the equality between the Casimirs (3.9), and the fact that if τ contains μ then it also contains the mirror image $\tilde{\mu}$. This is established towards the end of next Section, where also some comment on the exceptional cases is made. $\diamond$

We end this Section by noting that for the representations on de Sitter space $\hat{C}_\chi$, $\hat{C}_\tau$, there is no exhaustivity result as the Langlands-Knapp-Zuckerman result [57], [58] for ERs cited above. Thus, it is not surprising that not all conformal representations can be realized on de Sitter space, or, in other words, that some conformal fields live only on the boundary of de Sitter space and can not propagate into the bulk.

5. Intertwining relations between conformal and de Sitter representations

5.1. Bulk-to-boundary intertwining relation

This Section contains our main results, explicating the relations between CFT and de Sitter representations as intertwining relations. We first give in this subsection the intertwining operator from the de Sitter to the CFT realization. The operator which we use is mapping a function on de Sitter space to its boundary value and was used in a restricted sense (explained below) in many papers, starting from [11]. Also for those cases our result is new since we use it as operator between exactly defined spaces, and most importantly that we give it the interpretation of an intertwining operator.

Theorem: *Let us define the operator:*

$$L_\chi^\tau : \hat{C}_\chi^\tau \longrightarrow C_\chi, \quad (5.1)$$

with the following action:

$$(L_\chi^\tau \phi)(x) = \lim_{|a| \rightarrow 0} |a|^{-\Delta} \Pi_\mu^\tau \phi(x, |a|) \quad (5.2)$$

where Π_μ^τ is the standard projection operator from the K -representation space U_τ to the M -representation space V_μ , which acts in the following way on the K -representation matrices:

$$\Pi_\mu^\tau \hat{D}^\tau(k) = D^\mu(m(k)) \Pi_\mu^\tau \hat{D}^\tau(k_f) \quad (5.3)$$

where we have used (4.4). Then L_χ^τ is an intertwining operator, i.e.:

$$L_\chi^\tau \circ \hat{T}^\tau(g) = T^\chi(g) \circ L_\chi^\tau, \quad \forall g \in G. \quad (5.4)$$

In addition, in (5.2) the operator Π_μ^τ acts in the following truncated way:

$$\Pi_\mu^\tau \hat{D}^\tau(k) = D^\mu(m(k)) \Pi_\mu^\tau \quad (5.5)$$

Proof: Applying the LHS side of (5.4) to ϕ we have:

$$\begin{aligned} (L_\chi^\tau \circ \hat{T}^\mu(g)\phi)(x) &= \lim_{|a| \rightarrow 0} |a|^{-\Delta} \Pi_\mu^\tau (\hat{T}^\mu(g)\phi)(x, |a|) = \\ &= \lim_{|a| \rightarrow 0} |a|^{-\Delta} D^\mu(m(k)) \Pi_\mu^\tau \hat{D}^\tau(k_f) \phi(x', |a'|) , \end{aligned} \quad (5.6a)$$

$$g^{-1} \tilde{n}_x a = \tilde{n}_{x'} a' k^{-1} \quad (5.6b)$$

Applying the RHS side of (5.4) to ϕ we have:

$$\begin{aligned} (T^\chi(g) \circ L_\chi^\tau \phi)(x) &= |a''|^{-\Delta} D^\mu(m) (L_\chi^\tau \phi)(x'') = \\ &= |a''|^{-\Delta} D^\mu(m) \lim_{|a| \rightarrow 0} |a|^{-\Delta} \Pi_\mu^\tau \phi(x'', |a|) , \end{aligned} \quad (5.7a)$$

$$g^{-1} \tilde{n}_x = \tilde{n}_{x''} m^{-1} a''^{-1} n^{-1} \quad (5.7b)$$

In view of the Bruhat decomposition it is enough to prove coincidence between (5.6a) and (5.7a) for $g = \tilde{n}_y \in \tilde{N}, \hat{a} \in A, \hat{m} \in M$, and some element $w \in K, w \notin M$, representing the nontrivial element of the restricted Weyl group $W(G, A)$, since this element transforms elements of $\tilde{N}$ into elements of N : $w \tilde{n}_y w = n_{y'}$ (and thus makes unnecessary the check $g = n_y$). Such an element is, e.g., $w = \text{diag}(1, \dots, 1, -1, -1, 1)$, i.e., rotation in the plane $(d, d+1)$. However, for simplicity we shall demonstrate only the case of odd d when we can take $w \rightarrow R$ since in this case the conformal inversion $R \doteq \text{diag}(-1, \dots, -1, 1)$ is an element of K , (or we should suppose that we work with G'). We have:

- $g = \tilde{n}_y$: then (5.6b) gives $\tilde{n}_y^{-1} \tilde{n}_x a = \tilde{n}_{x-y} a$, i.e., $x' = x - y, a' = a, k = \mathbf{1}$, and (noting $\hat{D}^\tau(\mathbf{1}) = \mathbf{1}_\tau$) (5.6a) becomes:

$$\lim_{|a| \rightarrow 0} |a|^{-\Delta} \Pi_\mu^\tau \phi(x - y, |a|) \quad (5.8)$$

while (5.7b) gives $\tilde{n}_y^{-1} \tilde{n}_x = \tilde{n}_{x-y}$, i.e., $x'' = x - y, m = \mathbf{1}, a'' = \mathbf{1}, n = \mathbf{1}$, and (noting $D^\mu(\mathbf{1}) = \mathbf{1}_\mu$) (5.7a) also becomes (5.8).

- $g = \hat{a}$: then (5.6b) gives $\hat{a}^{-1} \tilde{n}_x a = \tilde{n}_{\frac{x}{|\hat{a}|}} \hat{a}^{-1} a$, i.e., $x' = \frac{x}{|\hat{a}|}, a' = \hat{a}^{-1} a, k = \mathbf{1}$, and (5.6a) becomes:

$$\lim_{|a| \rightarrow 0} |a|^{-\Delta} \Pi_\mu^\tau \phi\left(\frac{x}{|\hat{a}|}, \frac{|a|}{|\hat{a}|}\right) = |\hat{a}|^{-\Delta} \lim_{|a| \rightarrow 0} |a|^{-\Delta} \Pi_\mu^\tau \phi\left(\frac{x}{|\hat{a}|}, |a|\right) \quad (5.9)$$

while (5.7b) gives $\hat{a}^{-1} \tilde{n}_x = \tilde{n}_{\frac{x}{|\hat{a}|}} \hat{a}^{-1}$, i.e., $x'' = \frac{x}{|\hat{a}|}, m = \mathbf{1}, a'' = \hat{a}, n = \mathbf{1}$, and (5.7a) also becomes (5.9).

- $g = \hat{m}$: then (5.6b) gives $\hat{m}^{-1} \tilde{n}_x a = \tilde{n}_{\hat{m}^{-1}x} a \hat{m}^{-1}$, i.e., $x'_i = m_{ij}^{-1} x_j$, $a' = a$, $k = m(k) = \hat{m}$, and (5.6a) becomes:

$$\lim_{|a| \rightarrow 0} |a|^{-\Delta} \Pi_\mu^\tau \hat{D}^\tau(\hat{m}) \phi(x', |a|) = D^\mu(\hat{m}) \lim_{|a| \rightarrow 0} |a|^{-\Delta} \Pi_\mu^\tau \phi(x', |a|) \quad (5.10)$$

while (5.7b) gives $\hat{m}^{-1} \tilde{n}_x = \tilde{n}_{\hat{m}^{-1}x} \hat{m}^{-1}$, i.e., $x'' = x'$, $m = \hat{m}$, $a'' = \mathbf{1}$, $n = \mathbf{1}$, and (5.7a) also becomes (5.10).

- $g = R$: then (5.6b) gives $R \tilde{n}_x a = \tilde{n}_{x'} a' k^{-1}$ with:

$$\begin{aligned} x' &= x(x, a) \equiv -\frac{1}{x^2 + |a|^2} x, \\ a' &= a(x, a)^{-1} a, \quad |a(x, a)| = x^2 + |a|^2, \\ k &= k(x, a) \equiv \begin{pmatrix} \frac{2}{x^2 + |a|^2} x_i x_j - \delta_{ij} & -\frac{2|a|}{x^2 + |a|^2} x_i & 0 \\ \frac{2|a|}{x^2 + |a|^2} x_j & \frac{x^2 - |a|^2}{x^2 + |a|^2} & 0 \\ 0 & 0 & 1 \end{pmatrix} \end{aligned} \quad (5.11)$$

Using (4.5) we note for later use $k(x, \mathbf{1}) = -{}^t k_x$, and also:

$$k(x, a) = m(k(x, a)) k(x, |a|)_f = r(x) k_{\frac{|a|}{x^2}} x \quad (5.12)$$

Here we first record (for $x \neq 0$):

$$\begin{aligned} \lim_{|a| \rightarrow 0} x(x, a) &= -\frac{1}{x^2} x \equiv Rx \\ \lim_{|a| \rightarrow 0} a(x, a) &= a(x), \quad |a(x)| = x^2, \\ \lim_{|a| \rightarrow 0} k(x, a) &= r(x) \end{aligned} \quad (5.13)$$

and (5.6a) becomes:

$$\begin{aligned} \lim_{|a| \rightarrow 0} |a|^{-\Delta} \Pi_\mu^\tau \hat{D}^\tau(k(x, a)) \phi(x(x, a), |a(x, a)|^{-1} |a|) &= \\ &= (x^2)^{-\Delta} D^\mu(r(x)) \lim_{|a| \rightarrow 0} |a|^{-\Delta} \Pi_\mu^\tau \phi(Rx, |a|) \end{aligned} \quad (5.14)$$

On the other hand (5.7b) gives (also for $x \neq 0$): $R \tilde{n}_x = \tilde{n}_{x''} m^{-1} a''^{-1} n^{-1}$ with $x'' = Rx$, $m = r(x)$, $a'' = a(x)$, using the notation introduced in (5.13). Thus, (5.7a) also becomes (5.14).

This finishes the Proof of the intertwining property. On the way we have proved also (5.5). Indeed, though we have started with (5.3) in the generic f-la (5.6a) for the LHS of the intertwining property, in the four generating cases above we have $k_f = \mathbf{1}$, i.e., we could have started with (5.5) in (5.6a). ♠

5.2. Boundary-to-bulk intertwining relation

Now we look for the possible operator inverse to L_χ^τ which would restore a function on de Sitter space from its boundary value, as discussed in [11,14–33]. Again what is new here is that we define it as intertwining operator between exactly defined spaces in a more general setting. Moreover, we shall construct the operator just from the condition that it is an intertwining integral operator. Indeed, let us have the operator:

$$\tilde{L}_\chi^\tau : C_\chi \longrightarrow \hat{C}_\chi^\tau, \quad (5.15)$$

and try the following Ansatz:

$$\left(\tilde{L}_\chi^\tau f \right) (x, |a|) = \int K_\chi^\tau(x, |a|; x') f(x') dx' \quad (5.16)$$

where $K_\chi^\tau(x, |a|; x')$ is a linear operator acting from the space V_μ to the space U_τ , and let us suppose that $\tilde{L}_\chi^\tau$ is an intertwining operator, i.e.:

$$\hat{T}^\tau(g) \circ \tilde{L}_\chi^\tau = \tilde{L}_\chi^\tau \circ T^\chi(g), \quad \forall g \in G. \quad (5.17)$$

As in the Theorem above we apply (5.17) for $g = \tilde{n}_y, \hat{a}, \hat{m}, R$ and we use the same decompositions as above, so we can present things in a short fashion. Applying (5.17) for $g = \tilde{n}_y$ results in the fact that K_χ^τ depends only on the difference of the x arguments:

$$K_\chi^\tau(x, |a|; x') = K_\chi^\tau(x - x', |a|). \quad (5.18)$$

Applying (5.17) for $g = \hat{a}$ results in the fact that K_χ^τ is homogeneous in its arguments:

$$K_\chi^\tau(\sigma x, \sigma |a|) = \sigma^{\Delta-d} K_\chi^\tau(x, |a|), \quad \sigma \in \mathcal{C}, \sigma \neq 0 \quad (5.19)$$

Thus, we shall write:

$$K_\chi^\tau(x, |a|) = |a|^{\Delta-d} \hat{K}_\chi^\tau\left(\frac{x}{|a|}\right), \quad \hat{K}_\chi^\tau(y) \doteq K_\chi^\tau(y, 1) \quad (5.20)$$

Note now that (5.19) means, in particular, that $K_\chi^\tau(0, |a|)$ (if it exists) is fixed up to a constant matrix:

$$K_\chi^\tau(0, |a|) = |a|^{\Delta-d} K_\chi^\tau(0, 1) = |a|^{\Delta-d} \hat{K}_\chi^\tau(0). \quad (5.21)$$

Applying (5.17) for $g = \hat{m}$ results in the fact that K_χ^τ has the following covariance property:

$$\hat{D}^\tau(m) \hat{K}_\chi^\tau(x) = \hat{K}_\chi^\tau(mx) D^\mu(m), \quad \forall m \in M \quad (5.22)$$

The above means, in particular, that $\Pi_\mu^\tau \hat{K}_\chi^\tau(0)$ is M -invariant:

$$D^\mu(m) \Pi_\mu^\tau \hat{K}_\chi^\tau(0) = \Pi_\mu^\tau \hat{K}_\chi^\tau(0) D^\mu(m), \quad \forall m \in M \quad (5.23)$$

which then, by Schur's Lemma, means that $\Pi_\mu^\tau \hat{K}_\chi^\tau(0) = \sigma \mathbf{1}_\mu$, $0 \neq \sigma \in \mathcal{C}$. Thus, we have: $\hat{K}_\chi^\tau(0) = \sigma \Pi_\tau^\mu$, the latter being the canonical embedding operator from V_μ to U_τ , such that $\Pi_\mu^\tau \circ \Pi_\tau^\mu = \mathbf{1}_\mu$. Finally, applying (5.17) for $g = R$ means that:

$$\hat{D}^\tau(r(x)) \hat{D}^\tau(k_{\frac{|a|}{x^2}x}) \hat{K}_\chi^\tau(-\frac{x}{|a|} + \frac{x^2 + |a|^2}{|a|y^2}y) = \left(\frac{y^2}{x^2 + |a|^2}\right)^{d-\Delta} \hat{K}_\chi^\tau(\frac{x-y}{|a|}) D^\mu(r(y)) \quad (5.24)$$

where we have used the decomposition of $k(x, |a|)$ in (5.12). Now we set $y = x$ and we get:

$$\hat{D}^\tau(r(x)) \hat{D}^\tau(k_{\frac{|a|}{x^2}x}) \hat{K}_\chi^\tau(\frac{|a|}{x^2}x) = \left(\frac{x^2}{x^2 + |a|^2}\right)^{d-\Delta} \hat{K}_\chi^\tau(0) D^\mu(r(x)) \quad (5.25)$$

Using (5.22) for $x = 0$ we obtain (using also that $r(x)^2 = \mathbf{1}$):

$$\hat{D}^\tau(k_{\frac{|a|}{x^2}x}) \hat{K}_\chi^\tau(\frac{|a|}{x^2}x) = \left(\frac{x^2}{x^2 + |a|^2}\right)^{d-\Delta} \hat{K}_\chi^\tau(0) \quad (5.26)$$

Next, we make the change $\frac{|a|}{x^2}x \rightarrow x$ and we get:

$$\hat{K}_\chi^\tau(x) = N_\chi^\tau \left(\frac{1}{x^2 + 1}\right)^{d-\Delta} \hat{D}^\tau(k_{-x}) \Pi_\tau^\mu \quad (5.27)$$

or finally:

$$K_\chi^\tau(x, |a|) = N_\chi^\tau \left(\frac{|a|}{x^2 + |a|^2}\right)^{d-\Delta} \hat{D}^\tau(k_{-\frac{x}{|a|}}) \Pi_\tau^\mu \quad (5.28)$$

where N_χ^τ is arbitrary for the moment and should be fixed from the requirement that $\tilde{L}_\chi^\tau$ is inverse to L_χ^τ (when the latter is true).

The above operator exists for arbitrary representations τ of $K = SO(d+1)$ which contain the representation μ of $M = SO(d)$. We use the standard $SO(p)$ representation parametrization: $[\ell_1, \dots, \ell_{\tilde{p}}]$, ($\tilde{p} \equiv [\frac{p}{2}]$), where all ℓ_j are simultaneously integer or half-integer, all are positive except for p even when ℓ_1 can also be negative, and they are ordered: $|\ell_1| \leq \ell_2 \leq \dots \leq \ell_{\tilde{p}}$. The condition that $\tau = [\ell'_1, \dots, \ell'_{\hat{d}}]$, ($\hat{d} \equiv [\frac{d+1}{2}]$), contains $\mu = [\ell_1, \dots, \ell_{\tilde{d}}]$, ($\tilde{d} \equiv [\frac{d}{2}]$), explicitly is:

$$|\ell'_1| \leq \ell_1 \leq \dots \leq \ell_{\tilde{d}} \leq \ell'_{\hat{d}}, \quad d \text{ odd}, \quad \hat{d} = \tilde{d} + 1 \quad (5.29a)$$

$$-\ell'_1 \leq \ell_1 \leq \ell'_1 \leq \dots \leq \ell_{\tilde{d}} \leq \ell'_{\hat{d}}, \quad d \text{ even}, \quad \hat{d} = \tilde{d} \quad (5.29b)$$

If one is primarily concerned with the ERs $\chi = [\mu, \Delta]$ it is convenient to chose a 'minimal' representation $\tau(\mu)$ of $K = SO(d+1)$ containing μ . This depends on the parity of d . Thus, for μ as above, when d is odd we would choose:

$$\tau(\mu) = [\ell_1, \ell_1, \dots, \ell_{\tilde{d}}] \quad \text{or} \quad \tilde{\tau}(\mu) = [-\ell_1, \ell_1, \dots, \ell_{\tilde{d}}], \quad \mu \cong \tilde{\mu}, \quad (5.30)$$

while for *even* d we would choose:

$$\tau(\mu) = [|\ell_1|, \ell_2, \dots, \ell_{\tilde{d}}] = \tau(\tilde{\mu}) \cong \tilde{\tau}(\mu) = \tilde{\tau}(\tilde{\mu}), \quad \tilde{\mu} = [-\ell_1, \ell_2, \dots, \ell_{\tilde{d}}]. \quad (5.31)$$

Thus, in the odd d case for each μ we would choose between two K -irreps which are mirror images of one another, while in the even d case to each two mirror-image irreps of M we choose one and the same irrep of K .

The explicit formulae which appeared until now in the literature are actually in the cases in which $\tau = \tau(\mu)$, though there is no such interpretation as we have here. In such a restricted setting and from other considerations formula (5.28) for the scalar case (when both μ and $\tau = \tau(\mu)$ are scalar irreps) was given by Witten [11], while some other nonscalar cases were given in [11, 14, 17–19, 25–27]. Note that in (2.38) of [11] it is written for the conjugated conformal weight: $\Delta \rightarrow d - \Delta$, which in our language would mean to work with the representation $\tilde{\chi} = [\tilde{\mu}, d - \Delta]$ and to use:

$$K_{\tilde{\chi}}^{\tau}(x, |a|) = N_{\tilde{\chi}}^{\tau} \left(\frac{|a|}{x^2 + |a|^2} \right)^{\Delta} \hat{D}^{\tau}(k_{-\frac{x}{|a|}}) \Pi_{\tau}^{\tilde{\mu}} \quad (5.32)$$

5.3. Equivalence vs. partial equivalence

Either one of the representation equivalences established in the previous subsections means that the representations $\hat{C}_{\chi}^{\tau}$ and C_{χ} are *partially equivalent*. In order for them to be *equivalent* it is necessary and sufficient that the operators $\tilde{L}_{\chi}^{\tau}$, L_{χ}^{τ} are inverse to each other, i.e., the following relations should hold:

$$L_{\chi}^{\tau} \circ \tilde{L}_{\chi}^{\tau} = \mathbf{1}_{C_{\chi}}, \quad \tilde{L}_{\chi}^{\tau} \circ L_{\chi}^{\tau} = \mathbf{1}_{\hat{C}_{\chi}^{\tau}} \quad (5.33)$$

If (5.33) do not hold then at least one of the representations $\hat{C}_{\chi}^{\tau}$ and C_{χ} is topologically reducible.

For the first relation in (5.33) we have:

$$\begin{aligned} \left(L_{\chi}^{\tau} \circ \tilde{L}_{\chi}^{\tau} f \right)(x) &= \lim_{|a| \rightarrow 0} |a|^{-\Delta} \Pi_{\mu}^{\tau} \left(\tilde{L}_{\chi}^{\tau} f \right)(x, |a|) = \\ &= \lim_{|a| \rightarrow 0} |a|^{-\Delta} \Pi_{\mu}^{\tau} \int K_{\chi}^{\tau}(x - x', |a|) f(x') dx' \end{aligned} \quad (5.34)$$

For the above calculation we interchange the limit and the integration, and use the following result from [11] (there for $\Delta > d/2$):

$$\lim_{|a| \rightarrow 0} |a|^{\Delta-d} \left(\frac{|a|}{x^2 + |a|^2} \right)^{\Delta} \sim \delta(x) \quad (5.35)$$

substituting for our needs $\Delta \rightarrow d - \Delta$.

To obtain the proportionality constant in (5.35), and thus fix N_χ^τ , we first find the Fourier transform of $(x^2 + |a|^2)^{-\Delta}$:

$$\begin{aligned}
\int \frac{e^{-ip \cdot x}}{(x^2 + |a|^2)^\Delta} \frac{dx}{(2\pi)^{d/2}} &= \frac{1}{2^\Delta \Gamma(\Delta)} \int \int_0^\infty d\alpha \alpha^{\Delta-1} e^{-ip \cdot x - \frac{1}{2}\alpha(x^2 + |a|^2)} \frac{dx}{(2\pi)^{d/2}} = \\
&= \frac{1}{2^\Delta \Gamma(\Delta)} \int_0^\infty d\alpha \alpha^{\Delta-\frac{d}{2}-1} e^{-\frac{1}{2}\left(\frac{p^2}{\alpha} + \alpha|a|^2\right)} = \\
&= \frac{2}{2^\Delta \Gamma(\Delta)} \left(\frac{\sqrt{p^2}}{|a|}\right)^{\Delta-\frac{d}{2}} K_{\Delta-\frac{d}{2}}(|a|\sqrt{p^2})
\end{aligned} \tag{5.36}$$

where we have used f-la 3.47.9 of [65], involving the Bessel function K_ν . Note that for $|a| \rightarrow 0$ the above formula goes to formula (5.2) of [48] (with $\Delta = h + c$), in particular, the RHS of (5.36) goes to:

$$\frac{\Gamma(\frac{d}{2} - \Delta)}{2^\Delta \Gamma(\Delta)} \left(\frac{p^2}{2}\right)^{\Delta-\frac{d}{2}}, \quad \frac{d}{2} - \Delta \notin \mathbb{Z}_- \tag{5.37}$$

For (5.37) one uses the relation:

$$\lim_{\beta \rightarrow 0} 2 \beta^\nu K_\nu(\beta) = 2^\nu \Gamma(\nu), \quad \nu \notin \mathbb{Z}_- \tag{5.38}$$

Now we can find the necessary limit in (5.35) (again using (5.38)):

$$\begin{aligned}
\lim_{|a| \rightarrow 0} |a|^{\Delta-d} \left(\frac{|a|}{x^2 + |a|^2}\right)^\Delta &= \lim_{|a| \rightarrow 0} \frac{2 |a|^{2\Delta-d}}{2^\Delta \Gamma(\Delta)} \int \left(\frac{\sqrt{p^2}}{|a|}\right)^{\Delta-\frac{d}{2}} K_{\Delta-\frac{d}{2}}(|a|\sqrt{p^2}) \frac{e^{ip \cdot x} dp}{(2\pi)^{d/2}} = \\
&= \lim_{|a| \rightarrow 0} \frac{2}{2^\Delta \Gamma(\Delta)} \int \left(|a|\sqrt{p^2}\right)^{\Delta-\frac{d}{2}} K_{\Delta-\frac{d}{2}}(|a|\sqrt{p^2}) \frac{e^{ip \cdot x} dp}{(2\pi)^{d/2}} = \\
&= \frac{\Gamma(\Delta - \frac{d}{2})}{2^{\frac{d}{2}} \Gamma(\Delta)} \int \frac{e^{ip \cdot x} dp}{(2\pi)^{d/2}} = \\
&= \frac{\pi^{\frac{d}{2}} \Gamma(\Delta - \frac{d}{2})}{\Gamma(\Delta)} \delta(x)
\end{aligned} \tag{5.39}$$

where we have used:

$$\delta(x) = \int \frac{e^{ip \cdot x} dp}{(2\pi)^d}. \tag{5.40}$$

In the scalar case prompted by (5.39) (with $\Delta \rightarrow d - \Delta$) we choose the constant in (5.36) as:

$$N_\chi^\tau = \frac{\Gamma(d - \Delta)}{\pi^{\frac{d}{2}} \Gamma(\frac{d}{2} - \Delta)}, \quad \Delta \neq d + k, \quad k \in \mathbb{Z}_+ \text{ for } d \text{ odd}. \tag{5.41}$$

This is the choice made in [16] (for $\Delta \rightarrow d - \Delta$) from other considerations. In the general case we choose N_χ^τ as:

$$\begin{aligned}
N_\chi^\tau &= N_0 \frac{\Gamma(\hat{d} - \Delta)}{\Gamma(\frac{d}{2} - \Delta)} \prod_{k=1}^{\tilde{d}} (m_k + \frac{d}{2} - \Delta) \\
m_k &\doteq |\ell_k + k - 1 + \delta_d|, \quad k = 1, \dots, \tilde{d} \\
\tilde{d} &\equiv \left[\frac{d}{2}\right], \quad \hat{d} \equiv \left[\frac{d+1}{2}\right], \quad \delta_d = \frac{d}{2} - \tilde{d} = \begin{cases} 0 & d \text{ even} \\ \frac{1}{2} & d \text{ odd} \end{cases}
\end{aligned} \tag{5.42}$$

where N_0 is a constant independent of Δ having no poles or zeroes for any χ . Note that for d odd N_χ^τ has poles and thus is not defined in the following cases:

$$\Delta = \frac{d+1}{2} + p, \quad p \in \mathbb{Z}_+, \quad p \neq m_k - \frac{1}{2}, \quad k = 1, \dots, \tilde{d}, \quad d \text{ odd}, \tag{5.43}$$

(which for the scalar case coincide with the exclusion conditions in (5.41)). We note also the zero cases:

$$N_\chi^\tau = 0, \quad \text{for} \quad \begin{cases} \Delta = m_k + \frac{d}{2}, \quad k = 1, \dots, \tilde{d}, & d \text{ even} \\ \Delta = m_k + \frac{d}{2} \neq \frac{d+1}{2} + p, \quad k = 1, \dots, \tilde{d}, \quad p \in \mathbb{Z}_+, & d \text{ odd} \\ \Delta = \frac{d}{2} + p, \quad p \in \mathbb{Z}_+, & d \text{ odd} \end{cases} \tag{5.44}$$

Further we need also the following fact which for simplicity we write for d odd (or for $G \rightarrow G'$):

$$\lim_{|a| \rightarrow 0} k_{-\frac{x}{|a|}} = r(x) R \tag{5.45}$$

With the choice (5.41) (or (5.42)) and using (5.39) and (5.45) the last line of (5.34) gives $f(x)$ thus establishing the first relation in (5.33) for generic values of Δ (i.e., when N_χ^τ is finite and nonzero).

As a *Corollary* we recover the fact [11] that for generic values of Δ we can restore a function on de Sitter space from its boundary value on R^d . Indeed, suppose we have:

$$\phi(x, |a|) = \int K_\chi^\tau(x - x', |a|) f(x') dx', \quad \phi \in \hat{C}_\chi^\tau, \quad f \in C_\chi \tag{5.46}$$

Then we have for the boundary value:

$$\begin{aligned}
\psi_0(x) &\doteq (L_\chi^\tau \phi)(x) = \lim_{|a| \rightarrow 0} |a|^{-\Delta} \Pi_\mu^\tau \phi(x, |a|) = \\
&= \lim_{|a| \rightarrow 0} |a|^{-\Delta} \Pi_\mu^\tau \int K_\chi^\tau(x - x', |a|) f(x') dx' = f(x)
\end{aligned} \tag{5.47}$$

Now we can prove the second relation in (5.33):

$$\begin{aligned}
(\tilde{L}_\chi^\tau \circ L_\chi^\tau \phi)(x, |a|) &= \int K_\chi^\tau(x - x', |a|) (L_\chi^\tau \phi)(x') dx' = \\
&= \int K_\chi^\tau(x - x', |a|) \lim_{|a'| \rightarrow 0} |a'|^{-\Delta} \Pi_\mu^\tau \phi(x', |a'|) = \\
&= \int K_\chi^\tau(x - x', |a|) \psi_0(x') dx' = \phi(x, |a|)
\end{aligned} \tag{5.48}$$

where in the last line we used (5.47).

• Thus, we have found that the partially equivalent representations $\hat{C}_\chi^\tau$ and C_χ ($\chi = [\mu, \Delta]$) are equivalent iff Δ is not in the excluded ranges given in (5.43), (5.44).

This result may be used for the conjugate situation $\chi \rightarrow \tilde{\chi}$. The constant then is:

$$N_{\tilde{\chi}}^\tau = \tilde{N}_0 \frac{\Gamma(\Delta - \tilde{d})}{\Gamma(\Delta - \frac{d}{2})} \prod_{k=1}^{\tilde{d}} (m_k - \frac{d}{2} + \Delta) \tag{5.49}$$

where $\tilde{N}_0$ is a constant independent of Δ having no poles or zeroes for any χ . The cases when $N_{\tilde{\chi}}^\tau$ is not defined are:

$$\Delta = \tilde{d} - p, \quad p \in \mathbb{Z}_+, \quad p \neq m_k - \frac{1}{2}, \quad k = 1, \dots, \tilde{d}, \quad d \text{ odd}, \tag{5.50}$$

while the zero cases are:

$$N_{\tilde{\chi}}^\tau = 0, \quad \text{for} \quad \begin{cases} \Delta = \frac{d}{2} - m_k, & k = 1, \dots, \tilde{d}, & d \text{ even} \\ \Delta = \frac{d}{2} - m_k \neq \frac{d+1}{2} + p, & k = 1, \dots, \tilde{d}, p \in \mathbb{Z}_+, & d \text{ odd} \\ \Delta = \frac{d}{2} - p, & p \in \mathbb{Z}_+, & d \text{ odd} \end{cases} \tag{5.51}$$

• Thus, we find that the partially equivalent representations $\hat{C}_{\tilde{\chi}}^\tau$ and $C_{\tilde{\chi}}$ ($\tilde{\chi} = [\tilde{\mu}, d - \Delta]$) are equivalent iff Δ is not in the excluded ranges given in (5.50), (5.51).

5.4. Further intertwining relations

We start by recording the second limit of the bulk functions (which we mentioned towards the end of Section 4). We take as in the previous subsection $\psi_0 \in C_\chi$, $\chi = [\mu, \Delta]$, and $\phi \in \hat{C}_\chi^\tau$ expressed through ψ_0 by (5.46), (5.47). We set and calculate:

$$\begin{aligned}
\phi_0(x) &\doteq \lim_{|a| \rightarrow 0} |a|^{\Delta-d} \Pi_\mu^\tau \phi(x, |a|) = \\
&= \lim_{|a| \rightarrow 0} |a|^{\Delta-d} \Pi_\mu^\tau \int K_\chi^\tau(x - x', |a|) \psi_0(x') dx' = \\
&= N_\chi^\tau \int \frac{dx'}{(x - x')^{2(d-\Delta)}} \Pi_\mu^\tau \hat{D}^\tau(r(x)R) \Pi_\tau^\mu \psi_0(x') = \\
&= N_\chi^\tau \int \frac{dx'}{(x - x')^{2(d-\Delta)}} D^{\tilde{\mu}}(r(x)) \psi_0(x') = \\
&= \frac{N_\chi^\tau}{\gamma_{\tilde{\chi}}} \int dx' G_{\tilde{\chi}}(x - x') \psi_0(x')
\end{aligned} \tag{5.52}$$

Since $\psi_0 \in C_\chi$ from the intertwining property of $G_{\tilde{\chi}}$ follows that $\phi_0 \in C_{\tilde{\chi}}$, $\tilde{\chi} = [\tilde{\mu}, d - \Delta]$. This is valid for generic representations - in the exceptional cases it may happen that $\phi_0 = 0$ (when ψ_0 is in the kernel of $G_{\tilde{\chi}}$) or that the asymptotic expansion contains logarithms, (for the latter cf., e.g., (7.45) of [48], [66]).

Thus, we have established (4.9) at generic representation points. The two different limits of the bulk field ϕ are given by the coupled fields ϕ_0 and ψ_0 (the latter we denote also by $\mathcal{O}$).

Formulae (5.52) mean that if we define:

$$\mathcal{A}_{\tilde{\chi}}^\tau : \hat{C}_\chi^\tau \longrightarrow C_{\tilde{\chi}} \quad (5.53)$$

with the action:

$$(\mathcal{A}_{\tilde{\chi}}^\tau \phi)(x) = \lim_{|a| \rightarrow 0} |a|^{\Delta-d} \Pi_\mu^\tau \phi(x, |a|) \quad (5.54)$$

then we can show as in Theorem the intertwining property:

$$\mathcal{A}_{\tilde{\chi}}^\tau \circ \hat{T}^\tau(g) = T^{\tilde{\chi}}(g) \circ \mathcal{A}_{\tilde{\chi}}^\tau, \quad \forall g \in G, \quad (5.55)$$

since in the Proof of the Theorem we have used only the fact of the existence of the limit. On the other hand $\mathcal{A}_{\tilde{\chi}}^\tau$ is equal to $L_{\tilde{\chi}}^\tau$ because of (4.9).

Next we note that the intertwining property (5.55) is fulfilled if we take the following as a defining relation:

$$\mathcal{A}'_{\tilde{\chi}}^\tau = G_{\tilde{\chi}} \circ L_\chi^\tau \quad (5.56)$$

Expectedly, we get the same result as in (5.52) (up to a multiplicative constant):

$$\begin{aligned} (\mathcal{A}'_{\tilde{\chi}} \phi)(x) &= (G_{\tilde{\chi}} \circ L_\chi^\tau \phi)(x) = \\ &= \int dx' G_{\tilde{\chi}}(x - x') (L_\chi^\tau \phi)(x') = \\ &= \int dx' G_{\tilde{\chi}}(x - x') \psi_0(x') \end{aligned} \quad (5.57)$$

Gathering everything we have in this subsection we obtain the following relation between three of the operators under consideration:

$$L_{\tilde{\chi}}^\tau = \frac{N_\chi^\tau}{\gamma_{\tilde{\chi}}} G_{\tilde{\chi}} \circ L_\chi^\tau \quad (5.58)$$

At generic points from this we can obtain a lot of interesting relations, e.g., applying $\tilde{L}_\chi^\tau$ from the right we get:

$$L_{\tilde{\chi}}^\tau \circ \tilde{L}_\chi^\tau = \frac{N_\chi^\tau}{\gamma_{\tilde{\chi}}} G_{\tilde{\chi}} \quad (5.59)$$

which is in fact (5.52). Applying G_χ from the left we get:

$$L_\chi^\tau = \frac{\gamma_{\tilde{\chi}}}{N_\chi^\tau} G_\chi \circ L_{\tilde{\chi}}^\tau \quad (5.60)$$

and making the change $\chi \longleftrightarrow \tilde{\chi}$ in the last equality we get:

$$L_{\tilde{\chi}}^\tau = \frac{\gamma_\chi}{N_{\tilde{\chi}}^\tau} G_{\tilde{\chi}} \circ L_\chi^\tau \quad (5.61)$$

Comparing (5.58), (5.61), we obtain the following relation between the normalization constants:

$$N_\chi^\tau N_{\tilde{\chi}}^\tau = \gamma_\chi \gamma_{\tilde{\chi}} = C \rho(\chi) \quad (5.62)$$

where $\rho(\chi)$ is the analytic continuation of the Plancherel measure for the Plancherel formula contribution of the principal series of unitary irreps of G , and the last equality was shown in [67]. (The constant C is independent of χ .) The Plancherel measure itself is given as follows [68], [69]:⁶

$$\rho(\chi) = \rho(\tilde{\chi}) = \rho_0(\mu) \left(\prod_{1 \leq i < j \leq \tilde{d}} (m_j^2 - m_i^2) \right) \frac{\Gamma(\delta_d + c) \Gamma(\delta_d - c)}{\Gamma(c) \Gamma(-c)} \prod_{k=1}^{\tilde{d}} (m_k^2 - c^2) \quad (5.63)$$

$$c \equiv \Delta - \frac{d}{2}, \quad \rho_0(\mu) = \begin{cases} 1 & d \text{ even} \\ \prod_{k=1}^{\tilde{d}} m_k & d \text{ odd} \end{cases}$$

(cf. (5.42)). In the scalar case, when $\ell_j = 0$ for all j , we have:

$$\rho(\chi) = C' \frac{\Gamma(\frac{d}{2} + c) \Gamma(\frac{d}{2} - c)}{\Gamma(c) \Gamma(-c)} = C' \frac{\Gamma(\Delta) \Gamma(d - \Delta)}{\Gamma(\Delta - \frac{d}{2}) \Gamma(\frac{d}{2} - \Delta)} \quad (5.64)$$

the constant C' being independent of Δ .

Another consequence of (5.62) is that the constants N_χ^τ do not depend on τ since the constants γ_χ do not. Thus, we drop their superscript τ .

The above relation between these constants may be obtained also if we use the reconstruction of the field ϕ from ϕ_0 :

$$\phi(x, |a|) = \int K_{\tilde{\chi}}^\tau(x - x', |a|) \phi_0(x') dx' = \left(\tilde{L}_{\tilde{\chi}}^\tau \phi_0 \right)(x, |a|) \quad (5.65)$$

⁶ The principal series is obtained for c pure imaginary.

Proceeding as in (5.52) we have:

$$\begin{aligned}
\psi_0(x) &= (L_\chi^\tau \phi)(x) = \frac{N_{\tilde{\chi}}}{\gamma_\chi} \int dx' G_\chi(x-x') \phi_0(x') = \\
&= \frac{N_\chi N_{\tilde{\chi}}}{\gamma_\chi \gamma_{\tilde{\chi}}} \int dx' dx'' G_\chi(x-x') G_{\tilde{\chi}}(x'-x'') \psi_0(x'') = \\
&= \frac{N_\chi N_{\tilde{\chi}}}{\gamma_\chi \gamma_{\tilde{\chi}}} \psi_0(x)
\end{aligned} \tag{5.66}$$

where we have substituted (5.52) in the last but one line, and used (3.10) in the last line.

Another interesting relation includes a convolution of a K -kernel and a G -kernel. For this we apply $\tilde{L}_\chi^\tau$ to (5.59) from the left and changing $\chi \longleftrightarrow \tilde{\chi}$ we get (denoting $x_{ij} \equiv x_i - x_j$):

$$\tilde{L}_\chi^\tau \circ G_\chi = \frac{\gamma_\chi}{N_{\tilde{\chi}}} \tilde{L}_{\tilde{\chi}}^\tau \tag{5.67}$$

which applied to $f \in C_{\tilde{\chi}}$ gives:

$$\begin{aligned}
(\tilde{L}_\chi^\tau \circ G_\chi f)(x_1, |a|) &= \int dx_2 K_\chi^\tau(x_{12}, |a|) (G_\chi f)(x_2) = \\
&= N_\chi \gamma_\chi \int dx_2 dx_3 \left(\frac{|a|}{x_{12}^2 + |a|^2} \right)^{d-\Delta} \left(\frac{1}{x_{23}^2} \right)^\Delta \hat{D}^\tau(k_{-\frac{x_{12}}{|a|}}) \Pi_\tau^\mu D^\mu(r(x_{23})) f(x_3) = \\
&= N_\chi \gamma_\chi \int dx_2 dx_3 \left(\frac{|a|}{x_{12}^2 + |a|^2} \right)^{d-\Delta} \left(\frac{1}{x_{23}^2} \right)^\Delta \hat{D}^\tau\left(k_{-\frac{x_{12}}{|a|}} r(x_{23})\right) \Pi_\tau^{\tilde{\mu}} f(x_3) = \\
&= \gamma_\chi \int dx_3 \left(\frac{|a|}{x_{13}^2 + |a|^2} \right)^\Delta \hat{D}^\tau\left(k_{-\frac{x_{13}}{|a|}}\right) \Pi_\tau^{\tilde{\mu}} f(x_3) = \\
&= \frac{\gamma_\chi}{N_{\tilde{\chi}}} \int dx_3 K_{\tilde{\chi}}^\tau(x_{13}, |a|) f(x_3) = \frac{\gamma_\chi}{N_{\tilde{\chi}}} (\tilde{L}_{\tilde{\chi}}^\tau f)(x_1, |a|)
\end{aligned} \tag{5.68}$$

Finally, we notice that all operators that we have used may be found on the following commutative diagram:

$$\begin{array}{ccc}
\hat{C}_{\tilde{\chi}}^\tau & = & \hat{C}_\chi^\tau [\phi] \\
\\
L_{\tilde{\chi}}^\tau & \swarrow \nearrow & \tilde{L}_{\tilde{\chi}}^\tau \\
\\
C_{\tilde{\chi}}[\phi_0] & & C_\chi[\mathcal{O}, \psi_0]
\end{array} \tag{5.69}$$

6. Comments and outlook

From the point of view of calculating the conformal correlators [11,14–33] of the conformal field $\mathcal{O} \in C_\chi$, ($\chi = [\mu, \Delta]$, $\Delta > \frac{d}{2}$) from the de Sitter configuration described by the field $\phi \in \hat{C}_\chi^\tau$, the conditions (5.50), (5.50) are sufficient. If one needs to consider some of these exceptional Δ -values one has to take another choice for $N_{\tilde{\chi}}$ (e.g., like the one in [16] for $\Delta = \frac{d}{2}$), and thus obtain a nonzero operator $\tilde{L}_{\tilde{\chi}}^\tau$. But also then $\tilde{L}_{\tilde{\chi}}^\tau$ will not be inverse to $L_{\tilde{\chi}}^\tau$. Of course, one should repeat separately the calculations starting from the analogues of (5.36). Then one may conclude that the asymptotic behaviour of ϕ is logarithmic, cf., e.g., [16,22]. We do not discuss this anymore here, but we should note that on the conformal side logarithmic behaviour occurs also when one is renormalizing intertwining operators acting between reducible ERs, cf., e.g., [48], [51].

From the point of view of the direct correspondence between the conformal field $\mathcal{O} \in C_\chi$, and the bulk field $\phi \in \tilde{C}_\chi^\tau$ there are a lot of excluded points in the region directly interesting for the applications $\Delta > \frac{d}{2}$, cf. (5.43), (5.44). In particular, this means that some conformal representations can not be extended from the boundary to the bulk, cf. [7], [9], for discussion and earlier references. We do not consider this further here since this question should be discussed taking into account the subrepresentation structure of the reducible representations occurring at the excluded points, cf. [66].

Naturally, one should take into account also the supersymmetry aspects of the correspondence. There was a lot of work on this already, e.g., [5,6,8,12,13,34,36–38,40,42–47]. However, in order to implement our approach we shall need also results from [70–73] on the representations of the conformal superalgebra $su(2, 2/N)$ and supergroup $SU(2, 2/N)$. In particular, we shall use the classification of all positive energy unitary irreducible representations of $su(2, 2/N)$ (including explicit parametrization in terms of representation characteristics) [71], [72], the explicit parametrization of the superfield content of the massless UIRs of $su(2, 2/N)$ [71], the elementary representations of $SU(2, 2/N)$ and intertwining operators between ERs [73].

Acknowledgments. The author is thankful to the Deutsche Forschungsgemeinschaft (DFG) for the financial support as Guest Professor at ASI (TU Clausthal) and to the Director of ASI Prof. Dr. H.-D. Doebner for the hospitality. The author was partially supported by BNFR under contract Φ -643.

References

- [1] J. Maldacena, Adv. Theor. Math. Phys. **2** (1998) 231 (hep-th/971120).
- [2] M. Flato and C. Fronsdal, J. Math. Phys. **22** (1981) 1100.
- [3] E. Angelopoulos, M. Flato, C. Fronsdal and D. Sternheimer, Phys. Rev. **D23** (1981) 1278.
- [4] C. Fronsdal, Phys. Rev. **D26** (1982) 1988.
- [5] H. Nicolai and E. Sezgin, Phys. Lett. **143B** (1984) 103.
- [6] M. Gunaydin, P. van Nieuwenhuizen and N.P. Warner, Nucl. Phys. **B255** (1985) 63.
- [7] S. Ferrara and C. Fronsdal, Class. Quant. Grav. **15** (1998) 2153 (hep-th/9712239).
- [8] M. Gunaydin and D. Minic, Nucl. Phys. **B523** (1998) 145 (hep-th/9802047).
- [9] S. Ferrara and C. Fronsdal, Phys. Lett. **433B** (1998) 19 (hep-th/9802126).
- [10] S.S. Gubser, I.R. Klebanov and A.M. Polyakov, Phys. Lett. **428B** (1998) 105 (hep-th/9802109).
- [11] E. Witten, Adv. Theor. Math. Phys. **2** (1998) 253 (hep-th/9802150).
- [12] H.J. Kim, L.J. Romans and P. van Nieuwenhuizen, Phys. Rev. **D32** (1985) 389.
- [13] M. Gunaydin and N. Marcus, Class. Quant. Grav. **2** (1985) L11.
- [14] M. Henningson and K. Sfetsos, Phys. Lett. **431B** (1998) 63 (hep-th/9803251).
- [15] W. Mück and K.S. Viswanathan, Phys. Rev. **D58** (1998) 041901 (hep-th/9804035).
- [16] D.Z. Freedman, S.D. Mathur, A. Matusis and L. Rastelli, MIT-CTP-2727, hep-th/9804058.
- [17] H. Liu and A.A. Tseytlin, Nucl. Phys. **B533** (1998) 88 (hep-th/9804083).
- [18] G. Chalmers, H. Nastase, K. Schalm and R. Siebelink, ITP-SB-98-36, hep-th/9805105.
- [19] W. Mück and K.S. Viswanathan, Phys. Rev. **D58** (1998) 106006 (hep-th/9805145).
- [20] A.M. Ghezelbash, K. Kaviani, S. Parvizi and A.H. Fatollahi, Phys. Lett. **435B** (1998) 291 (hep-th/9805162).
- [21] S. Lee, S. Minwalla, M. Rangamani and N. Seiberg, PUPT-1796, hep-th/9806074.
- [22] A.M. Ghezelbash, M. Khorrami and A. Aghamohammadi, hep-th/9807034.
- [23] H. Liu and A.A. Tseytlin, IMPERIAL-TP-97-98-060, hep-th/9807097.
- [24] D.Z. Freedman, S.D. Mathur, A. Matusis and L. Rastelli, MIT-CTP-2770, hep-th/9808006.
- [25] S. Corley, hep-th/9808184.
- [26] A. Volovich, hep-th/9809009.
- [27] W.S. l'Yi, HEP-CNU-9810, hep-th/9809132; HEP-CNU-9812, hep-th/9811097.
- [28] E. D'Hoker and D.Z. Freedman, hep-th/9809179.
- [29] D. Berenstein, R. Corrado, W. Fischler and J. Maldacena, UTTG-05-98, hep-th/9809188.
- [30] G. Bonelli, SISSA-114-98-EP, hep-th/9810194.
- [31] M. Bianchi and S. Kovacs, ROM2F-98-37, hep-th/9811060.

- [32] H. Liu, hep-th/9811152.
- [33] E. D'Hoker and D.Z. Freedman, UCLA/98/TEP/34, MIT-CTP-2795, hep-th/9811257.
- [34] S. Ferrara, C. Fronsdal and A. Zaffaroni Nucl. Phys. **B532** (1998) 153 (hep-th/9802203).
- [35] M. Flato and C. Fronsdal, Lett. Math. Phys. **44** (1998) 249 (hep-th/9803013).
- [36] Y. Oz and J. Terning, Nucl. Phys. **B532** (1998) 163 (hep-th/9803167).
- [37] L. Andrianopoli and S. Ferrara, Phys. Lett. **430** (1998) 248 (hep-th/9803171).
- [38] S. Ferrara, M.A. Lledo and A. Zaffaroni, Phys. Rev. **D58** (1998) 105029 (hep-th/9805082).
- [39] V. Balasubramanian, P. Kraus and A. Lawrence, HUTP-98-A028, (hep-th/9805171).
- [40] M. Gunaydin, D. Minic and M. Zagermann, Nucl. Phys. **B534** (1998) 96 (hep-th/9806042).
- [41] S. Ferrara and C. Fronsdal, CERN-TH-98-186, hep-th/9806072.
- [42] S. Ferrara and A. Zaffaroni, CERN-TH-98-229, hep-th/9807090.
- [43] L. Andrianopoli and S. Ferrara, CERN-TH-98-234, hep-th/9807150.
- [44] M.J. O'Loughlin and S. Randjbar-Daemi, ICTP preprint IC/98/100, hep-th/9807208.
- [45] S. Ferrara, M. Porrati, A. Zaffaroni, CERN-TH-98-330, hep-th/9810063.
- [46] M. Gunaydin, D. Minic and M. Zagermann, PSU-TH-205, hep-th/9810226.
- [47] L. Andrianopoli and S. Ferrara, CERN-TH-98-388, hep-th/9812067.
- [48] V.K. Dobrev, G. Mack, V.B. Petkova, S.G. Petrova and I.T. Todorov, *Harmonic Analysis on the n - Dimensional Lorentz Group and Its Applications to Conformal Quantum Field Theory*, Lecture Notes in Physics, Vol. 63 (Springer, 1977).
- [49] K. Koller, Comm. Math. Phys. **40** (1975) 15.
- [50] A.W. Knap, *Representation Theory of Semisimple Groups (An Overview Based on Examples)*, (Princeton Univ. Press, 1986).
- [51] V.K. Dobrev and V.B. Petkova, Rep. Math. Phys. **13** (1978) 233.
- [52] V.K. Dobrev, Rep. Math. Phys. **25** (1988) 159.
- [53] I.T. Todorov, M.C. Mintchev and V.B. Petkova, *Conformal Invariance in Quantum Field Theory*, (Pisa, Sc. Norm. Sup., 1978).
- [54] G.M. Sotkov and D.Ts. Stoyanov, J. Phys. **A13** (1980) 2807.
- [55] B. Binengar, C. Fronsdal and W. Heidenreich, J. Math. Phys. **24** (1983) 2828.
- [56] V.B. Petkova and G.M. Sotkov, Lett. Math. Phys. **8** (1984) 217.
- [57] R.P. Langlands, On the classification of irreducible representations of real algebraic groups, Mimeographed notes, Princeton (1973).
- [58] A.W. Knap and G.J. Zuckerman, in: Lecture Notes in Math. Vol. 587 (Springer, Berlin, 1977) pp. 138-159.; Ann. Math. **116** (1982) 389-501.
- [59] A.M. Polyakov, JETP Lett. **12** (1970) 381.
- [60] S. Ferrara, A.F. Grillo, R. Gatto and G. Parisi, Lett. Nuovo Cim. **4** (1972) 115.

- [61] A.A. Migdal, 4-dimensional soluble models of conformal field theory, preprint, Landau Inst. Theor. Phys., Chernogolovka (1972).
- [62] E.S. Fradkin and M.Ya. Palchik, Nucl. Phys. **B99** (1975) 317.
- [63] Harish-Chandra, Acta Math. **113** (1965) 241; ibid. **116** (1966) 1.
- [64] W. Schmid, Rice University Studies, **56** (1970) 99;
R. Hotta, J. Math. Soc. Japan, **23** (1971) 384.
- [65] I.S. Gradshteyn and I.M. Ryzhik, *Tables of Integrals, Series and Products* (Academic Press, NY, 1965).
- [66] V.K. Dobrev, in preparation.
- [67] A.W. Knap, Ann. Math. **93** (1971) 489.
- [68] M.A. Naimark, Trudy Moskov. Mat. Obshch. **8** (1959) 121; English translation: Am. Math. Soc. Transl. **36** (1964) 101.
- [69] T. Hirai, Proc. Jap. Acad. **42** (1965) 323.
- [70] V.K. Dobrev and V.B. Petkova, Lett. Math. Phys. **9** (1985) 287.
- [71] V.K. Dobrev and V.B. Petkova, Phys. Lett. **162B** (1985) 127.
- [72] V.K. Dobrev and V.B. Petkova, Proceedings, eds. A.O. Barut and H.D. Doebner, Lecture Notes in Physics, Vol. 261 (Springer-Verlag, Berlin, 1986) p. 291 and p. 300.
- [73] V.K. Dobrev and V.B. Petkova, Fortschr. d. Phys. **35** (1987) 537-572.